

DSH PSYCHOTROPIC MEDICATION

Operational Procedures

ANTIDEPRESSANT PROTOCOL:

NOTE: Please see the section of the DSH Psychotropic Medication Policy regarding a specific protocol for selective serotonin reuptake inhibitor (SSRI) antidepressants (Chapter 31).

I. Indications:

- A. The treated individual cooperates with treatment and monitoring.
- B. At least one of the following clinical indications is present and documented in the chart. Note that some indications are limited to specific antidepressants.

- 1. DSM major depressive disorder.
- 2. DSM dysthymic disorder.
- 3. DSM bipolar mood disorder, depressed phase.

[NOTE: Tricyclic antidepressants should be avoided due to risk of rapid cycling. SSRI antidepressants and bupropion appear the least likely to induce switches into mania. Also, recent data indicates that antidepressants may offer limited efficacy in the context of treating the depressed phase of bipolar or schizoaffective disorder while increasing overall instability and the rate of mood cycling.]

- 4. DSM complicated bereavement.
- 5. Depressive signs and symptoms occurring comorbidly with another psychiatric or medical disorder, including postpartum depression. (Note that if a treatable cause of secondary depression is known, then treatment of the primary disorder should be the focus of clinical efforts.)
- 6. DSM seasonal affective disorder in which phototherapy is not available or not preferred.
- 7. DSM panic disorder

[NOTE: Tricyclic antidepressants, monoamine oxidase inhibitors (MAOIs), and SSRI antidepressants have the best established efficacy.]

- 8. DSM obsessive-compulsive disorder (OCD).

[NOTE: SSRI antidepressants and the tricyclic, clomipramine have demonstrated efficacy. In general; however, clomipramine should rarely be used due to its greater risk of adverse effect profile.]

- 9. DSM post-traumatic stress disorder (PTSD).

DSH PSYCHOTROPIC MEDICATION

Operational Procedures

[NOTE: SSRI antidepressants have the best established data regarding efficacy, however, almost all of the antidepressants have anecdotal support for associated depressive and anxiety symptoms.]

10. DSM acute stress disorder (ASD). [Efficacy is less well established than for PTSD. The SSRI antidepressants have been most studied.]
11. DSM social phobia. [Studies have been limited primarily to the SSRI antidepressants.]
12. DSM generalized anxiety disorder (GAD). [The SSRI antidepressants have been the most studied, with most studies showing moderate efficacy.]
13. Chronic pain syndromes.
 - a. Amitriptyline, nortriptyline, desipramine, and duloxetine have demonstrated efficacy.
 - b. Several antidepressants also have demonstrated prophylactic efficacy for migraine headaches.
 - c. Although milnacipran is indicated for treatment of fibromyalgia, levomilnacipran SR has not been indicated by the U.S. Food and Drug Administration for treatment of fibromyalgia but would likely be effective.
14. Smoking cessation. [Bupropion under the trade name Zyban has been the best studied.]
15. Eating disorders. [SSRI antidepressants have shown efficacy in bulimia nervosa and in binge eating disorder.]
16. DSM impulse control disorders and paraphilic disorders. [SSRI antidepressants have shown modest efficacy in some studies.]
17. Premature ejaculation. [SSRI antidepressants have shown efficacy in delaying orgasm.]

II. Contraindications:

- A. Hypersensitivity to the medication or any of the components of the formulation.
- B. Present hypomania or mania.
- C. For tricyclic antidepressants, current heart block or narrow angle glaucoma.
- D. For bupropion, current seizures due to unstable seizure disorder or unstable electrolyte status. Bupropion also carries a risk of diversion and abuse in forensic populations, as when insufflated or injected it acts as an amphetamine-like drug.

DSH PSYCHOTROPIC MEDICATION

Operational Procedures

- E. For venlafaxine, desvenlafaxine, duloxetine, or levomilnacipran SR, current unstable hypertension.
- F. For MAOIs, pregnancy. [MAOIs cause severe retardation of fetal growth across pregnancy, often causing fetal death.]

III. Precautions (documented risk/benefit analysis supports use):

- A. Hepatic disease (e.g., liver impairment may increase plasma concentrations)
 - 1. Nefazodone (no longer actively marketed in the U.S.) may cause acute necrosis.
- B. Renal disease (e.g., may cause accumulation of more toxic hydroxy metabolites).
- C. Cardiac disease (e.g., tricyclic antidepressants delay cardiac conduction)
 - 1. Desipramine may cause arrhythmia and sudden death in children.
- D. Hypertension (e.g., venlafaxine, desvenlafaxine, duloxetine, and levomilnacipran SR may increase diastolic pressures).
- E. Poorly controlled seizure disorder or unstable electrolyte status.
- F. Pregnancy or breast-feeding
 - 1. SSRI antidepressants cause decreased growth in the third trimester and may cause post-partum irritability, insomnia, and poor feeding.
 - 2. Paroxetine may cause cardiovascular defects in 1 per 25 live births with exposure in the first trimester. Half of the malformations are ventricular septal defects. Teratogenic risks were not supported by some later meta-analyses. Paroxetine also has caused post-partum seizures in the neonate.
- G. Prostatic hypertrophy
 - 1. Tricyclic antidepressants have substantial anticholinergic properties, especially the tertiary tricyclics.
 - 2. Levomilnacipran SR also may cause urinary retention.
- H. Like the SSRI antidepressants, other antidepressants may induce transient suicidal ideation.
- I. Those with serotonergic activity may decrease platelet aggregation and increase the risk of bleeding.

DSH PSYCHOTROPIC MEDICATION

Operational Procedures

IV. Pre-treatment workup:

- A. Informed consent or alternative legal authorization is present.
- B. Baseline workup includes:
 - 1. ECG within 1 year.
 - 2. Hepatic function tests within 30 days.
- C. BUN and creatinine within 30 days.
- D. EEG within 30 days if positive active seizure history is present.

V. Monitoring:

- A. Semi-annually:
 - 1. ECG with tricyclics for first 6 months.
 - 2. LFTs for first year for nefazodone, venlafaxine, and duloxetine only.
 - 3. BUN and creatinine for first year.
- B. Annually:
 - 1. LFTs.
 - 2. BUN and creatinine.
 - 3. ECG.

VI. Dosing:

- A. **Typical initial doses for the more commonly used heterocyclic antidepressants, excluding the SSRI antidepressants, are as follows:**

MEDICATION	INITIAL DOSE	TITRATION	MAXIMUM DOSE
Amitriptyline	25 - 50 mg/day	25 - 50 mg Q-2 to 4 days	300 mg/day
Amoxapine	25 - 50 mg/day	25 - 50 mg Q-2 to 4 days	300 mg/day
Bupropion	37.5 mg BID or TID	37.5 - 75 mg Q-3 to 7 days	150 mg TID
Bupropion SR	75 mg BID	75 to 100 mg Q-1 week	200 mg BID
Bupropion XL	150 mg QAM	150 mg Q-1 week	450 mg QAM
Clomipramine	25 - 50 mg QHS	25 - 50 mg Q-2 to 4 days	300 mg/day
Desipramine	25 mg QAM	25 - 50 mg Q-2 to 4 days	300 mg/day
Desvenlafaxine	50 mg QD	50 mg QD after 2 – 4 weeks	100 mg/day
Doxepine	25 - 50 mg QHS	25 - 50 mg Q-2 to 4 days	300 mg/day

DSH PSYCHOTROPIC MEDICATION

Operational Procedures

Duloxetine	30 mg/day	30 mg Q-1 week	120 mg/day
Imipramine	25 - 50 mg QHS	25 - 50 mg Q-2 to 4 days	300 mg/day
MEDICATION	INITIAL DOSE	TITRATION	MAXIMUM DOSE
Levomilnacipran SR	20 mg/day	Increase to 40 mg QD after 2 days, then titrate by 40 mg Q-2 days	120 mg/day
Maprotiline	25 mg QAM	25 mg Q-2 to 4 days	225 mg/day
Mirtazapine	7.5 - 15 mg QHS	7.5 - 15 mg Q-1 week	45 mg/day
Nortriptyline	25 - 50 mg/day	25 - 50 mg Q-2 to 4 days	150 mg/day
Protriptyline	10 mg QAM	10 mg Q-2 to 4 days	60 mg/day
Trazodone	25 - 50 mg QHS	25 - 50 mg Q-2 to 4 days	600 mg/day
Venlafaxine	37.5 mg BID	75 mg Q-1 week	375 mg/day
Venlafaxine XR	75 mg QAM	75 mg Q-1 week	225 mg/day

* Doses greater than the cited maximum maintained for more than 15 days require Therapeutic Review Committee (TRC) or Medication Review Committee (MRC) consultation.

* Please see the DSH Psychotropic Medication Policy regarding a specific protocol for the SSRI antidepressants.

B. Typical dosing for the monoamine oxidase inhibitors (MAOIs) is as follows:

MEDICATION	INITIAL DOSE	TITRATION	MAXIMUM DOSE
Isocarboxazid	10 mg/day	10 mg Q-1 week	30 mg/day
Phenylzine	30 mg/day	30 mg Q-1 week	90 mg/day
Transdermal Selegiline	6 mg/day	3 mg Q-1 week	12 mg/day
Tranylcypromine	15 mg/day	15 mg Q-1 week	60 mg/day

* Use of transcutaneous selegiline at doses of 6 mg per day does not typically require dietary restriction, but may prohibit use of sympathomimetic medications. Higher doses require the same dietary and medication restrictions as the other MAOIs, i.e. ingestion of tyramine rich foods or use of sympathomimetic medications may cause hypertensive crisis. Doses greater than the maximums cited above used for more than 15 days require MRC or TRC consultation.

C. Among elderly and medically fragile individuals, the initial dosing of antidepressants should generally be decreased by 30 to 50%. Also, the rate of titration should be slowed, such that the dose increase interval is increased by a factor of 2 to 3.

D. For the tricyclic antidepressants, it is worth noting that nortriptyline has an established therapeutic range of 75 – 150 ng/mL.

1. Although the therapeutic range is not as clearly defined for the other tricyclic antidepressants, it is worth noting that antidepressant efficacy is unlikely at plasma concentrations of <150 ng/mL, while increasing side-effects without further benefit becomes likely at plasma concentrations above 450 ng/mL.

DSH PSYCHOTROPIC MEDICATION

Operational Procedures

2. Importantly, recent data suggest that unlike previously thought, tricyclic antidepressants are not superior to other antidepressants for treatment-resistant depression. Thus, they should rarely, if ever, be used at antidepressant doses due to their narrow therapeutic index and consequent lethal risk.
- E. Because the tricyclic antidepressants may have a therapeutic index as low as 6 - 8. It should be noted that all of the tricyclic antidepressants are metabolized by cytochrome P450 2D6. Thus, drugs or substances which potentially inhibit 2D6 (e.g., paroxetine and fluoxetine) may increase tricyclic concentrations to toxic or lethal levels.
- F. Tricyclic antidepressants should never be added to an existing treatment with monoamine oxidase inhibitors, as the acute reuptake blockade of norepinephrine and/or serotonin may result in lethal hypertensive crisis or serotonin syndrome. In rare refractory cases, tricyclic antidepressants and MAOIs can be initiated together or MAOIs can be cautiously added to a tricyclic antidepressant treatment, however, these combinations still carry significant risks.
- G. Medications which induce hepatic enzymes (e.g, carbamazepine) may cause loss of antidepressant efficacy due to declines in plasma concentrations. Adjustment of dose based on measurement of plasma concentrations may be required in such cases.

References:

- Albert, U., Aguglia, E., Maina, G. & Bogetto, F. 2002. Venlafaxine versus clomipramine in the treatment of obsessive-compulsive disorder: a preliminary single-blind, 12-week, controlled study. *J Clin Psychiatry*, 63, 1004-9. doi: 10.4088/jcp.v63n1108.
- Alembic Pharmaceuticals Limited 2019. Desvenlafaxine Package Insert. Vadodara (India).
- Alexanderson, B. 1972. Pharmacokinetics of nortriptyline in man after single and multiple oral doses: the predictability of steady-state plasma concentrations from single-dose plasma-level data. *Eur J Clin Pharmacol*, 4, 82-91.
- Amneal Pharmaceuticals LLC 2019. Levomilnacipran Package Insert Bridgewater, New Jersey.
- Amrein, R., Stabl, M., Henauer, S., Affolter, E. & Jonkanski, I. 1997. Efficacy and tolerability of moclobemide in comparison with placebo, tricyclic antidepressants, and selective serotonin reuptake inhibitors in elderly depressed patients: a clinical overview. *Can J Psychiatry*, 42, 1043-50.
- Apotex Corp 2019. Trazodone: Package Insert and Label Information.
- Auclair, A., Martel, J., Assié, M., Bardin, I., Heusler, P., Cussac, D., Marien, M., Newman-Tancredi, A., O'Connor, J. & Depoortere, R. 2013. Levomilnacipran (F2695), a

DSH PSYCHOTROPIC MEDICATION

Operational Procedures

norepinephrine-preferring SNRI: profile in vitro and in models of depression and anxiety. *Neuropharmacology*, 70, 338-347.

Aurobindo Pharma Limited 2019. Duloxetine Package Insert East Windsor, New Jersey.

Aurobindo Pharma Limited 2019. Venlafaxine Extended Release Package Insert East Windsor, New Jersey.

Aurobindo Pharma Limited 2019. Venlafaxine Package Insert East Windsor, New Jersey.

Balant-gorgia, A. E., Gex-Fabry, M. & Balant, I. P. 1991. Clinical pharmacokinetics of clomipramine. *Clin Pharmacokinet*, 20, 447-62.

Bandelow, B., Reitt, M., Röver, C., Michaelis, S., Görlich, Y. & Wedekind, D. 2015. Efficacy of treatments for anxiety disorders: a meta-analysis. *Int Clin Psychopharmacol*, 30, 183-92.

Benkert, O., Gründer, G., Wetzel, H. & Hackett, D. 1996. A randomized, double-blind comparison of a rapidly escalating dose of venlafaxine and imipramine in inpatients with major depression and melancholia. *Journal of psychiatric research*, 30, 441-451.

Benkert, O., Muller, M. & Szegedi, A. 2002. An overview of the clinical efficacy of mirtazapine. *Human Psychopharmacology: Clinical and Experimental*, 17, S23-S26.

Bighelli, I., Castellazzi, M., Cipriani, A., Girlanda, F., Guaiana, G., Koesters, M., Turrini, G., Furukawa, T. A. & Barbui, C. 2018. Antidepressants versus placebo for panic disorder in adults. *Cochrane Database Syst Rev*, 4, Cd010676.

Birkenhager, T. K., Van den broek, W. W., Mulder, P. G., Bruijn, J. A. & Moleman, P. 2004. Efficacy and tolerability of tranylcypromine versus phenelzine: a double-blind study in antidepressant-refractory depressed inpatients. *Journal of Clinical Psychiatry*, 65, 1505-10.

Blob, I. F., Sharoky, M., Campbell, B. J., Kemper, E. M., Gilmor, M. G., Vandenberg, C. M. & Azzaro, A. J. 2007. Effects of a tyramine-enriched meal on blood pressure response in healthy male volunteers treated with selegiline transdermal system 6 mg/24 hour. *CNS Spectr*, 12, 25-34.

Bluepoint Laboratories 2020. Bupropion Extended Release (XL) Package Insert Ridgewood, New Jersey.

Bonnet, U. 2003. Moclobemide: therapeutic use and clinical studies. *CNS Drug Rev*, 9, 97-140.

Brannan, S. K., Mallinckrodt, C. H., Brown, E. B., Wohlreich, M. M., Watkin, J. G. & Schatzberg, A. F. 2005. Duloxetine 60 mg once-daily in the treatment of painful physical symptoms in patients with major depressive disorder. *Journal of psychiatric research*, 39, 43-53.

Brueckle, M. S., Thomas, E. T., Seide, S. E., Pilz, M., Gonzalez-Gonzalez, A. I., Nguyen, T. S., Harder, S., Glasziou, P. P., Gerlach, F. M. & Muth, C. 2020. Adverse drug reactions

DSH PSYCHOTROPIC MEDICATION

Operational Procedures

associated with amitriptyline - protocol for a systematic multiple-indication review and meta-analysis. *Syst Rev*, 9, 59.

Charlier, C., Pinto, E., Ansseau, M. & Plomteux, G. 2000. Relationship between clinical effects, serum drug concentration, and concurrent drug interactions in depressed patients treated with citalopram, fluoxetine, clomipramine, paroxetine or venlafaxine. *Hum Psychopharmacol*, 15, 453-459.

Cheng, Q., Huang, J., Xu, L., Li, Y., Li, H., Shen, Y., Zheng, Q. & Li, L. 2020. Analysis of time-course, dose-effect, and influencing factors of antidepressants in the treatment of acute adult patients with major depression. *Int J Neuropsychopharmacol*, 23, 76-87.

Chiuccariello, L., Cooke, R. G., Miler, L., Levitan, R. D., Baker, G. B., Kish, S. J., Kolla, N. J., Rusjan, P. M., Houle, S., Wilson, A. A. & Meyer, J. H. 2015. Monoamine oxidase-A occupancy by moclobemide and phenelzine: implications for the development of monoamine oxidase inhibitors. *Int J Neuropsychopharmacol*, 19.

Cipriani, A., Furukawa, T. A., Salanti, G., Geddes, J. R., Higgins, J. P., Churchill, R., Watanabe, N., Nakagawa, A., Omori, I. M. & McGuire, H. 2009. Comparative efficacy and acceptability of 12 new-generation antidepressants: a multiple-treatments meta-analysis. *The lancet*, 373, 746-758.

Cipriani, A., Koesters, M., Furukawa, T. A., Nosè, M., Purgato, M., Omori, I. M., Trespidi, C. & Barbui, C. 2012. Duloxetine versus other anti-depressive agents for depression. *Cochrane Database of Systematic Reviews*.

Clarke, A., Brewer, F., Johnson, E. S., Mallard, N., Hartig, F., Taylor, S. & Corn, T. H. 2003. A new formulation of selegiline: improved bioavailability and selectivity for MAO-B inhibition. *J Neural Transm (Vienna)*, 110, 1241-55.

Constantino, J. L. & Fonseca, V. A. 2019. Pharmacokinetics of antidepressants in patients undergoing hemodialysis: a narrative literature review. *Braz J Psychiatry*, 41, 441-446.

Davidson, J. R., Giller, E. L., Zisook, S. & Overall, J. E. 1988. An efficacy study of isocarboxazid and placebo in depression, and its relationship to depressive nosology. *Arch Gen Psychiatry*, 45, 120-7.

Dawling, S., Lynn, K., Rosser, R. & Braithwaite, R. 1981. The pharmacokinetics of nortriptyline in patients with chronic renal failure. *Br J Clin Pharmacol*, 12, 39-45.

Deecher, D. C., Beyer, C. E., Johnston, G., Bray, J., Shah, S., Abou-Gharbia, M. & Andree, T. H. 2006. Desvenlafaxine succinate: a new serotonin and norepinephrine reuptake inhibitor. *Journal of pharmacology and experimental therapeutics*, 318, 657-665.

Derry, S., Wiffen, P. J., Aldington, D. & Moore, R. A. 2015. Nortriptyline for neuropathic pain in adults. *Cochrane Database Syst Rev*, 1, Cd011209.

DSH PSYCHOTROPIC MEDICATION

Operational Procedures

Deupree, J. D., Montgomery, M. D. & Bylund, D. B. 2007. Pharmacological properties of the active metabolites of the antidepressants desipramine and citalopram. *Eur J Pharmacol*, 576, 55-60.

Dr. Reddy's Laboratories INC 2020. Bupropion Extended Release (SR) Package Insert Princeton, New Jersey.

Evans, L. E., Bett, J. H., Cox, J. R., Dubois, J. P. & Van Hees, T. 1980. The bioavailability of oral and parenteral chlorimipramine (Anafranil). *Prog Neuropsychopharmacol*, 4, 293-302.

Faulkner, R. D., Senekjian, H. O. & Lee, C. S. 1984. Hemodialysis of doxepin and desmethyldoxepin in uremic patients. *Artif Organs*, 8, 151-5.

Gaynes, B. N., Rush, A. J., Trivedi, M. H., Wisniewski, S. R., Spencer, D. & Fava, M. 2008. The STAR* D study: treating depression in the real world. *Cleveland Clinic journal of medicine*, 75, 57-66.

Gex-Fabry, M., Balant-Gorgia, A. E. & Balant, L. P. 1999. Clomipramine concentration as a predictor of delayed response: a naturalistic study. *Eur J Clin Pharmacol*, 54, 895-902.

Gillman, P. K. 2007. Tricyclic antidepressant pharmacology and therapeutic drug interactions updated. *Br J Pharmacol*, 151, 737-48.

Goodnick, P. J. 2007. Seligiline transdermal system in depression. *Expert Opinion on Pharmacotherapy*, 8, 59-64.

Gupta, S. K., Shah, J. C. & Hwang, S. S. 1999. Pharmacokinetic and pharmacodynamic characterization of OROS and immediate-release amitriptyline. *Br J Clin Pharmacol*, 48, 71-8.

Guy, S. & Silke, B. 1990. The electrocardiogram as a tool for therapeutic monitoring: a critical analysis. *J Clin Psychiatry*, 51 Suppl B, 37-9.

Hayasaka, Y., Purgato, M., Magni, L. R., Ogawa, Y., Takeshima, N., Cipriani, A., Barbui, C., Leucht, S. & Furukawa, T. A. 2015. Dose equivalents of antidepressants: Evidence-based recommendations from randomized controlled trials. *J Affect Disord*, 180, 179-84.

Hearn, L., Moore, R. A., Derry, S., Wiffen, P. J. & Phillips, T. 2014. Desipramine for neuropathic pain in adults. *Cochrane Database Syst Rev*, 2014, Cd011003.

Heritage Pharmaceuticals INC 2020. Bupropion Package Insert East Brunswick, New Jersey.

Herrera, D., Mayet, L., Galindo, M. C. & Jung, H. 2000. Pharmacokinetics of a sustained-release dosage form of clomipramine. *J Clin Pharmacol*, 40, 1488-93.

Hicks, J. K., Sangkuhl, K., Swen, J. J., Ellingrod, V. L., muller, d. J., shimoda, k., bishop, j. R., kharasch, e. D., skaar, t. C., gaedigk, a., dunnenberger, h. M., klein, t. E., caudle, k. E.

DSH PSYCHOTROPIC MEDICATION

Operational Procedures

& stingl, J. C. 2017. Clinical pharmacogenetics implementation consortium guideline (CPIC) for CYP2D6 and CYP2C19 genotypes and dosing of tricyclic antidepressants: 2016 update. *Clin Pharmacol Ther*, 102, 37-44.

Hilliard, W. T., Barloon, L., Farley, P., Penn, J. V. & Koranek, A. 2013. Bupropion diversion and misuse in the correctional facility. *Journal of Correctional Health Care*, 19, 211-217.

Hirschfeld, R. M., Mallinckrodt, C., Lee, T. C. & Detke, M. J. 2005. Time course of depression-symptom improvement during treatment with duloxetine. *Depression and anxiety*, 21, 170-177.

Kelly, M. W. & Myers, C. W. 1990. Clomipramine: a tricyclic antidepressant effective in obsessive compulsive disorder. *The Annals of Pharmacotherapy*, 24, 739-44.

Kennedy, S. H. 1997. Continuation and maintenance treatments in major depression: the neglected role of monoamine oxidase inhibitors. *Journal of Psychiatry & Neuroscience*, 22, 127-31.

Koechlin, B. A., Schwartz, M. A. & Oberhaensli, W. E. 1962. Metabolism of C-14-iproniazid and C-14-isocarboxazid in man. *J Pharmacol Exp Ther*, 138, 11-20.

Kriston, L., Von wolff, A., Westphal, A., Hölzel, L. P. & Härter, M. 2014. Efficacy and acceptability of acute treatments for persistent depressive disorder: a network meta-analysis. *Depress Anxiety*, 31, 621-30.

Kvist, E. E., Al-shurbaji, A., dahl, M. L., Nordin, C., Alván, G. & Ståhle, L. 2001. Quantitative pharmacogenetics of nortriptyline: a novel approach. *Clin Pharmacokinet*, 40, 869-77.

Larsen, J. K., Bendsen, B. B. & Bech, P. 2011. Vitamin B6 treatment of oedema induced by mirtazapine and isocarboxazid. *Acta Psychiatrica Scandinavica*, 124, 76-7; discussion 77.

Lee, K. C. & Chen, J. J. 2007. Transdermal selegiline for the treatment of major depressive disorder. *Neuropsychiatr Dis Treat*, 3, 527-37.

Leinonen, E., Lillsunde, P., Laukkanen, V. & Ylitalo, P. 1991. Effects of carbamazepine on serum antidepressant concentrations in psychiatric patients. *J Clin Psychopharmacol*, 11, 313-8.

Leucht, S., Steimer, W., Kreuz, S., Abraham, D., Orsulak, P. J. & Kissling, W. 2001. Doxepin plasma concentrations: is there really a therapeutic range? *J Clin Psychopharmacol*, 21, 432-9.

LIEBERMAN, J. A., COOPER, T. B., SUCKOW, R. F., STEINBERG, H., BORENSTEIN, M., BRENNER, R. & KANE, J. M. 1985. Tricyclic antidepressant and metabolite levels in chronic renal failure. *Clin Pharmacol Ther*, 37, 301-7.

DSH PSYCHOTROPIC MEDICATION

Operational Procedures

LOPEZ-MUNOZ, F. & ALAMO, C. 2009. Monoaminergic neurotransmission: the history of the discovery of antidepressants from 1950s until today. *Curr Pharm Des*, 15, 1563-86.

MACALUSO, M. & PRESKORN, S. H. 2011. CYP 2D6 PM status and antidepressant response to nortriptyline and venlafaxine: is it more than just drug metabolism? *J Clin Psychopharmacol*, 31, 143-5.

MALCOLM, D. E., YU, P. H., BOWEN, R. C., O'DONOVAN, C., HAWKES, J. & HUSSEIN, M. 1994. Phenelzine reduces plasma vitamin B6. *J Psychiatry Neurosci*, 19, 332-4.

MALLINGER, A. G., EDWARDS, D. J., HIMMELHOCH, J. M., KNOPF, S. & EHLER, J. 1986. Pharmacokinetics of tranylcypromine in patients who are depressed: relationship to cardiovascular effects. *Clin Pharmacol Ther*, 40, 444-50.

MALLINGER, A. G., HIMMELHOCH, J. M., THASE, M. E., EDWARDS, D. J. & KNOPF, S. 1990. Plasma tranylcypromine: relationship to pharmacokinetic variables and clinical antidepressant actions. *J Clin Psychopharmacol*, 10, 176-83.

MAYERSOHN, M. & GUENTERT, T. W. 1995. Clinical pharmacokinetics of the monoamine oxidase-A inhibitor moclobemide. *Clinical Pharmacokinetics*, 29, 292-332.

MCINTYRE, R. S., GOMMOLL, C., CHEN, C. & RUTH, A. 2016. The efficacy of levomilnacipran ER across symptoms of major depressive disorder: a post hoc analysis of 5 randomized, double-blind, placebo-controlled trials. *CNS Spectr*, 21, 385-392.

MOORE, R. A., DERRY, S., ALDINGTON, D., COLE, P. & WIFFEN, P. J. 2015. Amitriptyline for neuropathic pain in adults. *Cochrane Database Syst Rev*, 2015, Cd008242.

MORGAN, M. H. & READ, A. E. 1972. Antidepressants and liver disease. *Gut*, 13, 697-701.

MÜLLER, M. J., DRAGICEVIC, A., FRIC, M., GAERTNER, I., GRASMÄDER, K., HÄRTTER, S., HERMANN, E., KUSS, H. J., LAUX, G., OEHL, W., RAO, M. L., ROLLMANN, N., WEIGMANN, H., WEBER-LABONTE, M. & HIEMKE, C. 2003. Therapeutic drug monitoring of tricyclic antidepressants: how does it work under clinical conditions? *Pharmacopsychiatry*, 36, 98-104.

NAGY, A. & JOHANSSON, R. 1975. Plasma levels of imipramine and desipramine in man after different routes of administration. *Naunyn Schmiedebergs Arch Pharmacol*, 290, 145-60.

OLESEN, O. V. & LINNET, K. 1997. Metabolism of the tricyclic antidepressant amitriptyline by cDNA-expressed human cytochrome P450 enzymes. *Pharmacology*, 55, 235-43.

PAPAKOSTAS, G. I. & FAVA, M. 2007. A meta-analysis of clinical trials comparing milnacipran, a serotonin–norepinephrine reuptake inhibitor, with a selective serotonin reuptake inhibitor for the treatment of major depressive disorder. *European neuropsychopharmacology*, 17, 32-36.

DSH PSYCHOTROPIC MEDICATION

Operational Procedures

PAPAKOSTAS, G. I., THASE, M. E., FAVA, M., NELSON, J. C. & SHELTON, R. C. 2007. Are antidepressant drugs that combine serotonergic and noradrenergic mechanisms of action more effective than the selective serotonin reuptake inhibitors in treating major depressive disorder? A meta-analysis of studies of newer agents. *Biol Psychiatry*, 62, 1217-27.

PATKAR, A. A., PAE, C. U. & MASAND, P. S. 2006. Transdermal selegiline: the new generation of monoamine oxidase inhibitors. *CNS Spectr*, 11, 363-75.

PATRONEVA, A., CONNOLLY, S. M., FATATO, P., PEDERSEN, R., JIANG, Q., PAUL, J., GUICO-PABIA, C., ISLER, J. A., BURCZYNSKI, M. E. & NICHOLS, A. I. 2008. An assessment of drug-drug interactions: the effect of desvenlafaxine and duloxetine on the pharmacokinetics of the CYP2D6 probe desipramine in healthy subjects. *Drug Metab Dispos*, 36, 2484-91.

PERRY, P. J., ZEILMANN, C. & ARNDT, S. 1994. Tricyclic antidepressant concentrations in plasma: an estimate of their sensitivity and specificity as a predictor of response. *J Clin Psychopharmacol*, 14, 230-40.

PIGOTT, T. A. & SEAY, S. M. 1999. A review of the efficacy of selective serotonin reuptake inhibitors in obsessive-compulsive disorder. *J Clin Psychiatry*, 60, 101-6. doi: 10.4088/jcp.v60n0206.

POYUROVSKY, M. 2010. Acute antipsychotic-induced akathisia revisited. *The British Journal of Psychiatry*, 196, 89-91.

Preskorn, S., Patroneva, A., Silman, H., Jiang, Q., Isler, J. A., Burczynski, M. E., Ahmed, S., Paul, J. & Nichols, A. I. 2009. Comparison of the pharmacokinetics of venlafaxine extended release and desvenlafaxine in extensive and poor cytochrome P450 2D6 metabolizers. *Journal of clinical psychopharmacology*, 29, 39-43.

Preskorn, S. H. 2010. Understanding outliers on the usual dose-response curve: Venlafaxine as a way to phenotype patients in terms of their CYP 2D6 status and why it matters. *Journal of Psychiatric Practice*, 16, 46-49.

Rasmussen, B. B., Nielsen, T. L. & Brøsen, K. 1998. Fluvoxamine inhibits the CYP2C19-catalysed metabolism of proguanil in vitro. *Eur J Clin Pharmacol*, 54, 735-40. doi: 10.1007/s002280050544.

Remedyrepack INC 2020. Mirtazapine Package Insert Indiana, Pennsylvania.

Rheker, J., Rief, W., Doering, B. K. & Winkler, A. 2018. Assessment of adverse events in clinical drug trials: Identifying amitriptyline's placebo- and baseline-controlled side effects. *Exp Clin Psychopharmacol*, 26, 320-326.

Rief, W., Nestoriuc, Y., Von lilienfeld-toal, A., Dogan, I., Schreiber, F., Hofmann, S. G., Barsky, A. J. & Avorn, J. 2009. Differences in adverse effect reporting in placebo groups in SSRI and tricyclic antidepressant trials: a systematic review and meta-analysis. *Drug Saf*, 32, 1041-56.

DSH PSYCHOTROPIC MEDICATION

Operational Procedures

Robinson, D. S., Cooper, T. B., Jindal, S. P., Corcella, J. & Lutz, T. 1985. Metabolism and pharmacokinetics of phenelzine: lack of evidence for acetylation pathway in humans. *Journal of Clinical Psychopharmacology*, 5, 333-7.

Rodriguez de la torre, B., Dreher, J., Malevany, I., Bagli, M., Kolbinger, M., Omran, H., Luderitz, B. & Rao, M. L. 2001. Serum levels and cardiovascular effects of tricyclic antidepressants and selective serotonin reuptake inhibitors in depressed patients. *Therapeutic Drug Monitoring*, 23, 435-40.

Sallee, F. R. & Pollock, B. G. 1990. Clinical pharmacokinetics of imipramine and desipramine. *Clin Pharmacokinet*, 18, 346-64.

Scheiner, D. A., Perucchini, D., Fink, D. & Betschart, C. 2015. [Interstitial cystitis/bladder pain syndrome (IC/BPS)]. *Praxis (Bern 1994)*, 104, 909-18.

Schoerlin, M. P., Horber, F. F., Frey, F. J. & Mayersohn, M. 1990. Disposition kinetics of moclobemide, a new MAO-A inhibitor, in subjects with impaired renal function. *J Clin Pharmacol*, 30, 272-84.

Shulman, K. L., Herrmann, N. & Walker, S. E. 2013. Current place of monoamine oxidase inhibitors in the treatment of depression. *CNS Drugs*, 27, 789-97.

Soares, C. N., Endicott, J., Boucher, M., Fayyad, R. S. & Guico-Pabia, C. J. 2014. Predictors of functional response and remission with desvenlafaxine 50 mg/d in patients with major depressive disorder. *CNS Spectr*, 19, 519-527.

Spina, E., Avenoso, A., Campo, G. M., Caputi, A. P. & Perucca, E. 1995. The effect of carbamazepine on the 2-hydroxylation of desipramine. *Psychopharmacology (Berl)*, 117, 413-6.

Spina, E., Avenoso, A., Campo, G. M., Caputi, A. P. & Perucca, E. 1996. Phenobarbital induces the 2-hydroxylation of desipramine. *Ther Drug Monit*, 18, 60-4.

Stahl, S. M. 2009. Mechanism of action of trazodone: a multifunctional drug. *CNS Spectr*, 14, 536-546.

Stahl, S. M., Grady, M. M., Moret, C. & Briley, M. 2005. SNRIs: the pharmacology, clinical efficacy, and tolerability in comparison with other classes of antidepressants. *CNS Spectr*, 10, 732-747.

Stahl, S. M., Pradko, J. F., Haight, B. R., Modell, J. G., Rockett, C. B. & Learned-Coughlin, S. 2004. A review of the neuropharmacology of bupropion, a dual norepinephrine and dopamine reuptake inhibitor. *Primary care companion to the Journal of clinical psychiatry*, 6, 159.

Stimmel, G. L., Dopheide, J. A. & Stahl, S. M. 1997. Mirtazapine: an antidepressant with noradrenergic and specific serotonergic effects. *Pharmacotherapy: The Journal of Human Pharmacology and Drug Therapy*, 17, 10-21.

DSH PSYCHOTROPIC MEDICATION

Operational Procedures

Stoeckel, K., Pfefen, J. P., Mayersohn, M., Schoerlin, M. P., Andressen, C., Ohnhaus, E. E., Frey, F. & Guentert, T. W. 1990. Absorption and disposition of moclobemide in patients with advanced age or reduced liver or kidney function. *Acta Psychiatr Scand Suppl*, 360, 94-7.

Tasset, J. J., Singh, S. & Pesce, A. J. 1985. Evaluation of amitriptyline pharmacokinetics during peritoneal dialysis. *Ther Drug Monit*, 7, 255-7.

Thase, M. E. 1999. Antidepressant treatment of the depressed patient with insomnia. *The Journal of clinical psychiatry*.

Thase, M. E., Trivedi, M. H. & Rush, A. J. 1995. MAOIs in the contemporary treatment of depression. *Neuropsychopharmacology*, 12, 185-219.

Trinkley, K. E. & Nahata, M. C. 2014. Medication management of irritable bowel syndrome. *Digestion*, 89, 253-67.

Tsoi, D. T.-Y., Porwal, M. & Webster, A. C. 2010. Efficacy and safety of bupropion for smoking cessation and reduction in schizophrenia: systematic review and meta-analysis. *The British Journal of Psychiatry*, 196, 346-353.

Tyrer, P., Gardner, M., Lambourn, J. & Whitford, M. 1980. Clinical and pharmacokinetic factors affecting response to phenelzine. *British Journal of Psychiatry*, 136, 359-65.

Ulrich, S., Ricken, R. & Adli, M. 2017. Tranylcypromine in mind (Part I): Review of pharmacology. *Eur Neuropsychopharmacol*, 27, 697-713.

Ulrich, S., Ricken, R., Buspavanich, P., Schlattmann, P. & Adli, M. 2020. Efficacy and adverse effects of tranylcypromine and tricyclic antidepressants in the treatment of depression: a systematic review and comprehensive meta-analysis. *J Clin Psychopharmacol*, 40, 63-74.

Undurraga, J. & Baldessarini, R. J. 2017. Direct comparison of tricyclic and serotonin-reuptake inhibitor antidepressants in randomized head-to-head trials in acute major depression: Systematic review and meta-analysis. *J Psychopharmacol*, 31, 1184-1189.

Vandel, S., Bertschy, G., Baumann, P., Bouquet, S., Bonin, B., Francois, T., Sechter, D. & Bizouard, P. 1995. Fluvoxamine and fluoxetine: interaction studies with amitriptyline, clomipramine and neuroleptics in phenotyped patients. *Pharmacol Res*, 31, 347-53.

Venkatakrishnan, K., Greenblatt, D. J., Von Moltke, L. L., Schmider, J., Harmatz, J. S. & Shader, R. I. 1998. Five distinct human cytochromes mediate amitriptyline N-demethylation in vitro: dominance of CYP 2C19 and 3A4. *Journal of Clinical Pharmacology*, 38, 112-21.

Von Ammon Cavanaugh, S. 1990. Drug-drug interactions of fluoxetine with tricyclics. *Psychosomatics*, 31, 273-6.

Wang, W. A., Qian, J. M. & Pan, G. Z. 2003. [Treatment of refractory irritable bowel syndrome with subclinical dosage of antidepressants]. *Zhongguo Yi Xue Ke Xue Yuan Xue Bao*, 25, 74-8.

DSH PSYCHOTROPIC MEDICATION

Operational Procedures

Yan, J. H., Hubbard, J. W., McKay, G., Korchinski, E. D. & Midha, K. K. 2002. Absolute bioavailability and stereoselective pharmacokinetics of doxepin. *Xenobiotica*, 32, 615-23.

Yue, Q. Y., Zhong, Z. H., Tybring, G., Dalén, P., Dahl, M. L., Bertilsson, L. & Sjöqvist, F. 1998. Pharmacokinetics of nortriptyline and its 10-hydroxy metabolite in Chinese subjects of different CYP2D6 genotypes. *Clin Pharmacol Ther*, 64, 384-90.

Ziegler, V. E., Biggs, J. T., Wylie, L. T., Rosen, S. H., Hawf, D. J. & Coryell, W. H. 1978. Doxepin kinetics. *Clin Pharmacol Ther*, 23, 573-9.

Zisook, S. 1984. Side effects of isocarboxazid. *J Clin Psychiatry*, 45, 53-8.